



Corrigendum

Corrigendum to “Percutaneous or surgical revascularization in patients with severe left main coronary artery disease in Latin America: A GRADE clinical practice guideline” [International Journal of Cardiology Volume 436, 1 October 2025, 133401]

Pablo Lamelas^{a,b,*}, Michel Pompeu Sá^{c,1}, Ariel Izcovich^d, Federico Bottaro^e, Matias Tisi Baña^f, Maria Ines Sosa Liprandi^g, Fernando Lanas^h, Omar Asdrúbal Vilca Mejiaⁱ, Mauricio Zuñiga Luna^j, Patricia Aubanel^k, Ana Munera^l, Juan Contreras Reyes^m, Rodrigo Bagurⁿ, Richard Whitlock^{b,o}, Hector Garcia Garcia^p, Mamas Mamas^q, Mauricio G. Cohen^r, Alejandro Ricalde^s, Alexandre Abizaid^t, Oscar Mendiz^a, Mario Araya^u, Ricardo Costa^v, Omar Santaera^w, Pedro Hidalgo^x, Tulio Caldonazo^{y,z}, Adrian Baranchuk^{aa}, Martín Alberto Ragusa^{ab}, Endorsed by the Sociedad Latino Americana de Cardiología Intervencionista (SOLACI), the Sociedad Interamericana de Cardiología (SIAC)

^a Fundación Favaloro, Buenos Aires, Argentina

^b Health Research Methods, Evidence, and Impact, McMaster University, Canada

^c Department of Cardiovascular Surgery, Pronto Socorro Cardiológico de Pernambuco (PROCAPE), University of Pernambuco (UPE), Brazil

^d Hospital Alemán, Buenos Aires, Argentina

^e Department of Internal Medicine, Hospital Británico, Buenos Aires, Argentina

^f Department of Medicine, Hospital Universitario Austral, Pilar, Buenos Aires, Argentina

^g Sanatorio Güemes, Buenos Aires, Argentina

^h Universidad de la Frontera, Temuco, Chile

ⁱ Department of Cardiovascular Surgery, Instituto de Coracao, Universidade de Sao Paulo, Brazil

^j Department of Interventional Cardiology, Angiografía de Occidente, Cali, Colombia

^k Hospital del Prado, Tijuana, Mexico

^l Clínica Rosario Tesoro, Cardioestudio, Medellín, Colombia

^m Department of Cardiac Surgery, Universidad de la Frontera, Temuco, Chile

ⁿ Western University, London, Ontario, Canada

^o Department of Surgery, McMaster University, Hamilton, Ontario, Canada

^p Washington Hospital Center, Washington, USA

^q Keele Cardiovascular Research Group, Keele University, Staffordshire, UK

^r Cleveland Clinic Florida, Weston, FL, USA

^s Hospital ABC, Ciudad de México, Mexico

^t Instituto do Coração, Sao Paulo, Brazil

^u Clínica Alemana, Universidad del Desarrollo, Santiago de Chile, Chile

^v Instituto Dante Pazzanese de Cardiología, Sao Paulo, Brazil

^w Clínica privada provincial de Merlo, Buenos Aires, Argentina

^x Policlínica Amado, Maracaibo, Venezuela

^y Department of Cardiothoracic Surgery, Jena University Hospital, Jena, Germany

^z Department of Cardiothoracic Surgery, Weil Cornell Medicine, New York, USA

^{aa} Queens University, Kingston, Ontario, Canada

^{ab} División Clínica Médica, Hospital Fernández, Buenos Aires, Argentina

*Corresponding author at: Fundación Favaloro, Buenos Aires, Argentina; Health Research Methods, Evidence, and Impact, McMaster

DOI of original article: <https://doi.org/10.1016/j.ijcard.2025.133401>.

* Corresponding author.

E-mail address: pablolamelas@hotmail.com (P. Lamelas).

¹ Co-first author.

<https://doi.org/10.1016/j.ijcard.2025.133520>

Available online 9 July 2025

0167-5273/© 2025 The Author(s). Published by Elsevier B.V. All rights are reserved, including those for text and data mining, AI training, and similar technologies.

University, Canada

The authors of the manuscript asked for the support of the Latin-American Association of Cardiac and Endovascular Surgery (LACES),

however no common ground on the composition of the panel and methodology group committee could be found and therefore the authors decided to develop the guideline without the support of LACES.